



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

portional to CM and BC . From C measure the distance CD —this third proportional. D is the required point.

For $BC^2 = CM \cdot CD$, or $BC^2 = (AB - BC)CD$. $\therefore CD \cdot AB = BC^2 + BC \cdot CD$.

$$CD \cdot AB + BC \cdot CD + CD^2 = BC^2 + 2BC \cdot CD + CD^2.$$

$$CD(AB + BC + CD) = (BC + CD)^2.$$

$$AD \cdot CD = BD^2.$$

Also solved by S. A. COREY, J. R. HITT, F. D. POSEY, M. E. GRAEBER, W. W. LANDIS, and G. W. GREENWOOD.

249. Proposed by W. W. BEMAN, The University of Michigan.

Given the distances of a point in the plane of a square from three of its vertices, to find the side of the square.

250. Proposed by W. W. BEMAN, The University of Michigan.

Given the distances of a point in the plane of an equilateral triangle from the vertices; to find the side of triangle. [Perkins' *Geometry*, Olney's *Geometry*.]

I. Solution by F. D. POSEY, San Mateo, Calif.

Consider the general case, viz: Given the distances of a point in the plane of a regular n -gon to three consecutive vertices, to find the side of the n -gon.

Let the vertices be A, B, C in order, say clockwise, and P the given point. Let $PA, PB, PC \equiv a, b, c$, respectively. Let $\angle ABP = \alpha$, $\angle PBC = \beta$, taking these clockwise if P be without the angle ABC , and counter-clockwise if P be within. Call the side of the n -gon, x .

There are now two cases: (1) P within the angle ABC of the n -gon, (2) P without this angle. In the first case $\alpha + \beta = \frac{n-2}{n}\pi$. $\therefore \cos \beta = -\cos \frac{2\pi}{n} \cos \alpha + \sin \frac{2\pi}{n} \sin \alpha$. $\therefore \sin \alpha = \operatorname{cosec} \frac{2\pi}{n} \cos \beta + \cot \frac{2\pi}{n} \cos \alpha$.

In the second case $\alpha + \beta = 2\pi - \frac{n-2}{n}\pi$. $\therefore \sin \alpha = -\operatorname{cosec} \frac{2\pi}{n} \cos \beta - \cot \frac{2\pi}{n} \cos \alpha$

Now $\cos \alpha = \frac{b^2 + x^2 - a^2}{2bx}$ (when c is between b and a we have $\cos(2\pi - \alpha) = \cos \alpha$), and $\cos \beta = \frac{b^2 + x^2 - c^2}{2bx}$ (when a is between b and c we have $\cos(2\pi - \beta) = \cos \beta$). In both cases (1) and (2), $\sin^2 \alpha + \cos^2 \alpha = (\operatorname{cosec} \frac{2\pi}{n} \cos \beta + \cot \frac{2\pi}{n} \cos \alpha)^2 + \cos^2 \alpha = 1$, which equation after substituting the above values for $\cos \alpha$ and $\cos \beta$ reduces to:

$$\begin{aligned} & \left[\left(\cot \frac{2\pi}{n} + \operatorname{cosec} \frac{2\pi}{n} \right)^2 + 1 \right] x^4 + 2 \left\{ \left(\cot \frac{2\pi}{n} + \operatorname{cosec} \frac{2\pi}{n} \right) \left[\cot \frac{2\pi}{n} (b^2 - a^2) \right. \right. \\ & \left. \left. + \operatorname{cosec} \frac{2\pi}{n} (b^2 - c^2) \right] - a^2 - b^2 \right\} x^2 + \left[\cot \frac{2\pi}{n} (b^2 - a^2) + \operatorname{cosec} \frac{2\pi}{n} (b^2 - c^2) \right]^2 \\ & \quad + (b^2 - a^2)^2 = 0. \end{aligned}$$